


I NEED TO STAY HOME IF...

I HAVE A FEVER	I AM VOMITTING	I HAVE DIARRHEA	I HAVE A RASH	I HAVE HEAD LICE	I HAVE AN EYE INFECTION	I HAVE BEEN IN THE HOSPITAL
						
TEMPERATURE OF 100 OR HIGHER	WITHIN THE PAST 24 HOURS	WITHIN THE PAST 24 HOURS	BODY RASH WITH ITCHING OR FEVER	ITCHY HEAD, ACTIVE HEAD LICE	REDNESS, ITCHING, AND/OR "CRUSTY" DRAINAGE FROM EYE	HOSPITAL STAY AND/OR ER VISIT

I AM READY TO GO BACK TO SCHOOL WHEN I AM...

FEVER FREE FOR 24 HOURS WITHOUT THE USE OF FEVER REDUCING MEDICATIONS (i.e. TYLENOL, MOTRIN)	FREE FROM VOMITTING FOR AT LEAST 2 SOLID MEALS	FREE FROM DIARRHEA FOR AT LEAST 24 HOURS	FREE FROM RASH ITCHING OR FEVER. I HAVE BEEN EVALUATED BY MY DOCTOR IF NEEDED.	TREATED WITH LICE TREATMENT AT HOME AND PROOF IS PROVIDED	EVALUATED BY MY DOCTOR AND HAVE NOTE TO RETURN TO SCHOOL.	RELEASE BY MY MEDICAL PROVIDER TO RETURN TO SCHOOL.
--	--	--	--	---	---	---

Note: You know your child best! Did they sleep well? Do they have enough energy to participate in the day? Did the pediatrician give you guidance on when to return

GUIDELINES:

- **Strep throat**
 - At least 24 hours after starting antibiotics and fever free 24 hours.
- **Flu**
 - Typically, 5 days after onset. See symptom guide chart
- **Ear infection**
 - At least 24 hours after starting antibiotics and fever free 24 hours.
- **RSV**
 - May be contagious for 3-8 days. Can return to school after being fever free 24 hours.
- **Croup**
 - May return 2-3 days post illness symptoms. Must be fever free, feel well enough to return.
- **Coxsackievirus**
 - Sores have dried and fever free 24 hours.
- **Chicken Pox**
 - When scabs dry